

Unique preduals of Banach spaces: Property (X) of the predual of $H^\infty(\mathbb{D})$

MARIO GUILLÉN, PEDRO TRADACETE

Departamento de Matemática Aplicada, Universitat Politècnica de València

mguisan3@upv.edu.es

Resumen. This work introduces the concept of uniqueness of predual, shows examples from the literature and proves that property (X) is a sufficient condition for being a unique predual. Lastly, we review the fact that the predual of $H^\infty(\mathbb{D})$ has property (X), providing proofs not found in the literature.

Palabras clave: Unique preduals; Banach spaces; Property (X); Space of bounded holomorphic functions.

References

- [1] E. Amar and A. Lederer (1971). Points exposés de la boule unité de $H^\infty(\mathbb{D})$. *Comptes Rendus de l'Académie des Sciences. Série A*, vol. 272, pp. 1449-1552.
- [2] , T. Ando (1978). On the predual of H^∞ . *Commentationes Mathematicae: Tomus Specialis in Honorem Ladislai Orlicz*, vol. 1, pp. 33-40.
- [3] G. Godefroy (1980). Nouvelles classes d'espaces de Banach à prédual unique. *Séminaire Analyse fonctionnelle (dit "Maurey-Schwartz")*, 1980, pp. 1-28
- [4] G. Godefroy. Existence and uniqueness of isometric preduals: A survey. *Banach space theory* (Proc., Iowa City, 1987). Ed. by Bor-Luh Lin. Vol. 85. 1989, pp. 131-193.
- [5] H. Pfitzner (2007). Separable L-embedded Banach spaces are unique preduals. *Bulletin of the London Mathematical Society* 39.
- [6] M. Guillén (2025). Unique preduals and free objects in Banach spaces (Master thesis). <https://riunet.upv.es/handle/10251/225022>

Agradecimientos. This project has been developed under the JAE Intro Scholarship from the Consejo Superior de Investigaciones Científicas (CSIC), supervised by Pedro Tradacete

Indicar la preferencia (subrayar la opción elegida): póster o charla.

Indicar la preferencia (subrayar la opción elegida): Lunes/Martes o Jueves/Viernes.